

February 1, 2008

hep-ph/9301294

A More-Effective Potential *

Kevin Cahill

*Department of Physics and Astronomy, University of New Mexico**Albuquerque, New Mexico 87131-1156 [†]**Theoretical Physics Group, Lawrence Berkeley Laboratory**Berkeley, California 94720*

Abstract

In theories with spontaneous symmetry breaking, the loop expansion of the effective potential is awkward. In such theories, the exact effective potential $V(\phi_c, T)$ is real and convex (as a function of the classical field ϕ_c), but its perturbative series can be complex with a real part that is concave. These flaws limit the utility of the effective potential, particularly in studies of the early universe. A generalization of the effective potential is available that is real, that has no obvious convexity problems, and that can be computed in perturbation theory. For the theory with classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$, this more-effective potential closely tracks the usual effective potential where the latter is real $|\phi_c| \geq \sigma/\sqrt{3}$ and naturally extends it to $\phi_c = 0$, revealing that the critical temperature at the one-loop level runs from $T_C \approx 1.81\sigma$ for $\lambda = 0.1$ to $T_C \approx 1.74\sigma$ for $\lambda = 1$.

*This work was supported by the Director, Office of Energy Research, Office of High-Energy and Nuclear Physics, Division of High-Energy Physics of the U.S. Department of Energy under contracts DE-AC03-76SF00098 and DE-FG04-84ER40166.

[†]Permanent address, e-mail: kevin@cahill.unm.edu

I Introduction

The effective potential was introduced by Heisenberg and Euler [1] and by Schwinger [2]. Goldstone, Salam, and Weinberg [3] and Jona-Lasinio [4] developed the effective potential and applied it to the problem of symmetry breaking [5]. Coleman and E. Weinberg used it to show that radiative corrections could break symmetries [6]. Linde [7] and S. Weinberg [8] later used it to obtain a lower bound on the mass of the Higgs boson. West and others have used it to study the breaking of supersymmetry [9].

The possibility that broken symmetries might be restored at high temperatures was raised by Kirzhnits and Linde [10] and confirmed by them [11], by Dolan and Jackiw [12] and by S. Weinberg [13] who with Bernard [14] introduced and developed the finite-temperature effective potential. Much current work on the early universe is based upon the finite-temperature effective potential [15].

Although the effective potential has a long history of successful applications to particle physics, it does not seem to be well suited to theories that exhibit spontaneous symmetry breaking. In such theories the loop expansion of the effective potential can be awkward [16]. While the exact effective potential is both real and convex (as a function of the mean values ϕ_c of the scalar fields) [17], its perturbative series in theories with spontaneously broken symmetry displays neither property. In the example provided by the symmetry-breaking classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$, the loop expansion of the finite-temperature effective potential is complex at all temperatures T for $|\phi_c| < \sigma/\sqrt{3}$, and its real part is concave at low temperatures and small λ for a similar range of $|\phi_c|$. In such theories the accuracy of the one-loop effective potential does not extend down to the small values of $|\phi_c|$ that are of interest in studies of the early universe.

Of these two issues, convexity and complexity, it is the complexity that is the more serious. The convexity of the exact effective potential (with an ultraviolet regulator in place) suggests that the effective potential may not be the ideal tool for studying systems with spontaneous symmetry breaking in quasi-equilibrium in finite regions of spacetime. The concavity of its renormalized loop expansion in turn suggests that the loop expansion may be an uncertain approximation to the effective potential for such systems. But the critical defect of the effective potential is the complexity of its loop expansion. For where the effective potential is complex, it is ambiguous as an approximation to a free-energy density — although it may be interpreted as a decay rate [18].

Because of these limitations of the perturbative effective potential, some physicists have turned to nonperturbative techniques. Chang [19] has invented a variational method called the gaussian effective potential, which Barnes and Ghandour [20] and Stevenson [21] have developed. Fukuda and Kyriakopoulos [22] have introduced a version of the effective potential that is well suited to lattice computations; O’Raifeartaigh, Wipf, and Yoneyama [23] have analyzed this potential. Ringwald and Wetterich [24] have suggested the use of block-spin techniques.

The goal of the present paper is to generalize the effective potential so that it can be applied simply and perturbatively to theories with spontaneous symmetry breaking. The usual effective potential is the Legendre transform of the Helmholtz free-energy density for the modified hamiltonian $H + \int j\phi d^3x$ in which j is an external source. For theories with potentials of positive curvature, $V''(\phi) \geq 0$, the probe $j\phi$ is optimal. But for theories in which $V''(\phi)$ takes on negative values, as it must when $V(\phi)$ has two minima, it is argued that the linear probe $j\phi$ be generalized to a quadratic polynomial $jP(\phi)$. This advice has been given in the past with varying degrees of obliqueness by Cornwall, Jackiw,

and Tomboulis [25], by Hawking and Moss [26], and by Lawrie [27]; but it has not been followed. When discussing theories with spontaneous symmetry breaking, most physicists either ignore the complexity of the usual effective potential [15] or work in a region of parameter space in which scalar loops can be ignored [28].

In what follows I shall discuss the case of a single scalar field ϕ interacting with an arbitrary renormalizable potential $V(\phi)$. If the curvature $V''(\phi)$ of the potential is positive, then the usual effective potential with a linear probe $j\phi$ is adequate. But if curvature $V''(\phi)$ of the potential is negative for some values of the field ϕ , then a quadratic polynomial $jP(\phi)$ should be used. If the potential $V(\phi)$ of indefinite curvature has a single minimum at ϕ_1 , then a suitable probe is $P(\phi) = (\phi - \phi_1)^2/2$. If the potential $V(\phi)$ of indefinite curvature has two minima separated by a local maximum at ϕ_0 , then I suggest using $P(\phi) = (\phi - \phi_0)^2/2$. Such quadratic probes $jP(\phi)$ lead to more-effective potentials that possess real loop expansions and have no obvious convexity problems.

For the classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$, which has a local maximum at $\phi_0 = 0$, the appropriate probe is $P(\phi) = \phi^2/2$. The resulting effective potential closely tracks the usual effective potential where the latter is real and naturally extends it down to $\phi_c = 0$. The reflection symmetry of the action is restored to the vacuum at a critical temperature T_C which runs from $T_C \approx 1.81\sigma$ for $\lambda = 0.1$ to $T_C \approx 1.74\sigma$ for $\lambda = 1$. The first few terms of the high-temperature expansion of the one-loop effective potential $V_1(\phi_c, T; P)$ with probe $P(\phi) = \phi^2/2$ are

$$V_1(\phi_c, T; P) = -\frac{\pi^2}{90}T^4 + \frac{\lambda}{24}(3\phi_c^2 - \sigma^2)T^2 - \frac{\lambda^{3/2}}{12\sqrt{2}\pi}(7|\phi_c|^3 - 3\sigma^2|\phi_c|)T. \quad (1)$$

The corresponding terms of the usual effective potential are

$$V_1(\phi_c, T) = -\frac{\pi^2}{90}T^4 + \frac{\lambda}{24}(3\phi_c^2 - \sigma^2)T^2 - \frac{\lambda^{3/2}}{12\pi}(3\phi_c^2 - \sigma^2)^{3/2}T. \quad (2)$$

These two expansions possess the same two leading terms, but they differ in the

third term, which is imaginary for $|\phi_c| < \sigma/3$ in the case of the usual effective potential. In the expansion of the more-effective potential $V_1(\phi_c, T; P)$, the term $\lambda^{3/2}\sigma^2|\phi_c|T$ occurs with a positive coefficient and may have astrophysical implications[29].

The traditional effective potential is discussed in Sections II–V in a pedagogical manner inspired by Weinberg [13]. The problems that can arise in theories that exhibit spontaneous symmetry breaking are illustrated in Sec. VI. The more-effective potential is introduced in Sec. VII. The meaning of generalized effective potentials is discussed in Sec. VIII. In Sec. IX the computation of the generalized effective potential is discussed for the case of a quartic polynomial $V(\phi)$ with two minima. This computation is carried out in detail for the potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$ in Sec. X.

II The Partition Function of a Free Field

One of the clearest descriptions of the finite-temperature effective potential is S. Weinberg’s operator formulation [13]. Stripped of fermions and gauge fields and reduced to a single scalar field, it will serve as the basis for the introductory sections of this paper.

A free, real scalar field ϕ of mass m is described by the hamiltonian

$$\begin{aligned} H &= \int \frac{1}{2} \left(\pi^2 + (\nabla\phi)^2 + m^2\phi^2 \right) d^3x \\ &= \sum_k \omega_k \left[a^\dagger(k)a(k) + \frac{1}{2} \right] \end{aligned} \tag{3}$$

where $\omega_k = \sqrt{k^2 + m^2}$. At temperature T its partition function is

$$Z(\beta) = \text{Tr } e^{-\beta H} \tag{4}$$

where $\beta = 1/T$ is the inverse temperature. By inserting a complete set of energy

eigenstates, one may find for the partition function $Z(\beta)$ the expression

$$\begin{aligned} Z(\beta) &= \prod_k e^{-\frac{\beta\omega_k}{2}} \sum_{n_k=0}^{\infty} e^{-\beta n_k \omega_k} \\ &= \prod_k \frac{e^{-\frac{\beta\omega_k}{2}}}{(1 - e^{-\beta\omega_k})} \end{aligned} \quad (5)$$

which is simpler as a logarithm,

$$\begin{aligned} -\log Z(\beta) &= \sum_k \frac{\beta\omega_k}{2} + \log(1 - e^{-\beta\omega_k}) \\ &= \int \frac{L^3 d^3k}{(2\pi)^3} \left[\frac{\beta\omega_k}{2} + \log(1 - e^{-\beta\omega_k}) \right] \end{aligned} \quad (6)$$

where L^3 is the volume of quantization.

III The Effective Potential

For a scalar field ϕ described by a hamiltonian H perturbed by an external current j , the Helmholtz free-energy density $A(j, T)$ is defined by the relation

$$e^{-\beta L^3 A(j, T)} = \text{Tr} e^{-\beta(H + j \int \phi(x) d^3x)}. \quad (7)$$

The free energy $A(j, T)$ is therefore proportional to the logarithm of the partition function $Z(\beta, j)$

$$A(j, T) = -\frac{\log Z(\beta, j)}{\beta L^3} \quad (8)$$

for the system described by the perturbed hamiltonian $H + j \int \phi(x) d^3x$.

The mean value of the field ϕ is a function of the current j

$$\phi_c(j, T) \equiv \langle \phi \rangle_j = \frac{\text{Tr} \phi(x) e^{-\beta(H + j \int \phi(x) d^3x)}}{\text{Tr} e^{-\beta(H + j \int \phi(x) d^3x)}} \quad (9)$$

and is a derivative of the Helmholtz potential

$$\phi_c(j, T) = \frac{\partial A(j, T)}{\partial j}. \quad (10)$$

The finite-temperature effective potential $V(\phi_c, T)$ is defined [10–12] as a Legendre transform of the Helmholtz potential

$$V(\phi_c, T) \equiv A(j, T) - j \frac{\partial A(j, T)}{\partial j} = A(j, T) - j\phi_c \quad (11)$$

expressed as a function of the “classical field” ϕ_c

$$V(\phi_c, T) = A(j(\phi_c, T), T) - j(\phi_c, T)\phi_c \quad (12)$$

rather than of the current j . The effective potential may be thought of as a Gibbs free-energy density. It is obviously real.

The utility of the effective potential derives from its ability at its minima to represent the unperturbed system. For from eqs.(10) and (11), it follows that the derivative of the effective potential with respect to the classical field ϕ_c is proportional to the external current j

$$\frac{\partial V(\phi_c, T)}{\partial \phi_c} = \frac{\partial j}{\partial \phi_c} \left(\frac{\partial A(j, T)}{\partial j} - \phi_c \right) - j = -j. \quad (13)$$

Thus the current j must vanish at the stationary points of $V(\phi_c, T)$,

$$0 = \frac{\partial V(\phi_c, T)}{\partial \phi_c} = -j. \quad (14)$$

At zero temperature, the minimum value of the effective potential is the energy density of the ground state of the system.

Since it is through the factor $\exp(-\beta j \int \phi(x) d^3x)$ that the current j influences the mean value ϕ_c , the relationship between the current j and the mean value ϕ_c is inverse. Thus both the derivative of the mean value ϕ_c with respect to the current j and that of j with respect to ϕ_c are negative or zero: $\partial \phi_c / \partial j \leq 0$ and $\partial j / \partial \phi_c \leq 0$. So by differentiating the formula (13) with respect to ϕ_c , one sees that the effective potential has a nonnegative second derivative,

$$\frac{\partial^2 V(\phi_c, T)}{\partial \phi_c^2} = -\frac{\partial j}{\partial \phi_c} \geq 0. \quad (15)$$

The effective potential is therefore formally convex [17] as a function of the field ϕ_c .

IV The Effective Potential for a Free Field

In the case of a free scalar field, one may implement these definitions exactly.

The unitary transformation

$$U(j) = e^{-i \int (j/m^2) \pi(x) d^3x} \quad (16)$$

displaces the field $\phi(x)$

$$U^\dagger \phi(x) U = \phi(x) + \frac{j}{m^2} \quad (17)$$

and so relates the perturbed hamiltonian to the unperturbed one:

$$U^\dagger H U = H + \int j \phi(x) d^3x + \frac{L^3 j^2}{2m^2}. \quad (18)$$

Thus since traces are invariant under unitary transformations, the Helmholtz potential $A(j, T)$ for the free field

$$\begin{aligned} e^{-\beta L^3 \left(A + \frac{j^2}{2m^2} \right)} &= \text{Tr} e^{-\beta \left(H + \int j \phi d^3x + \frac{L^3 j^2}{2m^2} \right)} \\ &= \text{Tr} U^\dagger e^{-\beta H} U \\ &= \text{Tr} e^{-\beta H} = Z(\beta) \end{aligned} \quad (19)$$

is related to the logarithm (6) of the partition function of the unperturbed system by the equation

$$A(j, T) = -\frac{\log Z(\beta)}{\beta L^3} - \frac{j^2}{2m^2}. \quad (20)$$

The effect of the linear perturbation $j \int \phi(x) d^3x$ is to displace the field by j/m^2 , as shown by eq.(17). So the mean value ϕ_c is

$$\phi_c = \frac{\partial A(j, T)}{\partial j} = -\frac{j}{m^2}. \quad (21)$$

One may also evaluate ϕ_c directly. Since the mean value $\langle\phi\rangle$ for the unperturbed theory is $\phi_c(0, T) = 0$, that of the perturbed theory is simply

$$\begin{aligned}\phi_c(j, T) &= \frac{\text{Tr } \phi(x) U^\dagger e^{-\beta H} U}{\text{Tr } U^\dagger e^{-\beta H} U} \\ &= \frac{\text{Tr } U \phi(x) U^\dagger e^{-\beta H}}{\text{Tr } e^{-\beta H}} \\ &= -\frac{j}{m^2}.\end{aligned}\tag{22}$$

It follows now from the definition (11), from eqs.(20–22), and from the formula (6) for the partition function $Z(\beta)$ that the exact finite-temperature effective potential for the free scalar field of mass m is

$$\begin{aligned}V(\phi_c, T) &= A(j, T) - j\phi_c \\ &= \frac{1}{2}m^2\phi_c^2 - \frac{\log Z(\beta)}{\beta L^3} \\ &= \frac{1}{2}m^2\phi_c^2 + \int \frac{d^3k}{(2\pi)^3} \left[\frac{\omega_k}{2} + \frac{\log(1 - e^{-\beta\omega_k})}{\beta} \right]\end{aligned}\tag{23}$$

with $\omega_k = \sqrt{k^2 + m^2}$. The effective potential $V(\phi_c, T)$ is real and convex. At its absolute minimum $\phi_c = 0$, the external current $j = -m^2\phi_c$ vanishes. In the limit $\beta \rightarrow \infty$, the potential $V(\phi_c, T)$ becomes the exact zero-temperature effective potential

$$V(\phi_c, 0) = \frac{1}{2}m^2\phi_c^2 + \int \frac{d^3k}{(2\pi)^3} \frac{\omega_k}{2}.\tag{24}$$

V The One-Loop Effective Potential

For a scalar field interacting with itself through a potential $V(\phi)$, the effect of the perturbing current j is to replace $V(\phi)$ by

$$V_j(\phi) = V(\phi) + j\phi.\tag{25}$$

The absolute minimum $\bar{\phi}$ of this altered potential is a root of the equation

$$0 = \frac{\partial V_j(\phi)}{\partial \phi} = \frac{\partial V(\phi)}{\partial \phi} + j. \quad (26)$$

To obtain the one-loop approximation to the Helmholtz potential, we replace the altered potential $V_j(\phi)$ in the definition (7) of $A(j, T)$ by its Taylor-series expansion about the absolute minimum $\bar{\phi}$:

$$V_j(\phi) \approx V_j(\bar{\phi}) + \frac{1}{2} \frac{\partial^2 V_j(\bar{\phi})}{\partial \phi^2} (\phi - \bar{\phi})^2. \quad (27)$$

To zeroth order in $\hbar$, the minimum $\bar{\phi}$ is the mean value ϕ_c of the scalar field ϕ as defined by eq.(9). The truncated series (27)

$$V_j(\phi) \approx V(\bar{\phi}) + j\bar{\phi} + \frac{1}{2} \frac{\partial^2 V(\bar{\phi})}{\partial \phi^2} (\phi - \bar{\phi})^2 \quad (28)$$

describes a free scalar field of mass

$$m = \sqrt{V''(\bar{\phi})}. \quad (29)$$

The quantity $V''(\bar{\phi})$ is positive because $\bar{\phi}$ is a minimum of $V_j(\phi)$.

One may now express the Helmholtz potential $A(j, T)$ in terms of the kinetic energy $K = \int (1/2)\pi(x)^2 d^3x$ as

$$\begin{aligned} e^{-\beta L^3 A(j, T)} &= \text{Tr} e^{-\beta(H + j \int \phi(x) d^3x)} \\ &= \text{Tr} e^{-\beta(K + \int V_j(x) d^3x)} \\ &\approx e^{-\beta L^3 (V(\bar{\phi}) + j\bar{\phi})} \text{Tr} e^{-\beta [K + \int (1/2)m^2 (\phi(x) - \bar{\phi})^2 d^3x]}. \end{aligned} \quad (30)$$

So by using the unitary operator

$$U = e^{i \int \bar{\phi} \pi(x) d^3x} \quad (31)$$

which displaces the field $\phi(x)$ to $U^\dagger \phi(x) U = \phi(x) - \bar{\phi}$, one may write $A(j, T)$ approximately as

$$e^{-\beta L^3 A(j, T)} \approx e^{-\beta L^3 (V(\bar{\phi}) + j\bar{\phi})} \text{Tr} [U^\dagger e^{-\beta H_0} U]$$

$$\begin{aligned}
&\approx e^{-\beta L^3(V(\bar{\phi})+j\bar{\phi})} \text{Tr}[e^{-\beta H_0}] \\
&\approx e^{-\beta L^3(V(\bar{\phi})+j\bar{\phi})} Z(\beta)
\end{aligned} \tag{32}$$

where $Z(\beta)$ is the exact partition function (6) for the free scalar field of mass m . Thus at the one-loop level, the Helmholtz potential is

$$A_1(j, T) = V(\bar{\phi}) + j\bar{\phi} - \frac{\log Z(\beta)}{\beta L^3}. \tag{33}$$

The mean value ϕ_c and the mean $\bar{\phi}$ differ only by terms of order $\hbar$, which are due to the quantum fluctuations induced by the kinetic energy K . Specifically it follows from eqs.(10) and (26) that this difference is

$$\phi_c = \bar{\phi} - \frac{\partial}{\partial j} \frac{\log Z(\beta)}{\beta L^3}. \tag{34}$$

Thus by the extremal condition (26), the altered potential changes only by quantities that are of second order in $\hbar$ as $\bar{\phi}$ is replaced by ϕ_c ,

$$V(\bar{\phi}) + j\bar{\phi} = V(\phi_c) + j\phi_c + \mathcal{O}(\hbar^2). \tag{35}$$

We may therefore write the Helmholtz potential to first order in $\hbar$ in terms of the mean value ϕ_c of the field ϕ

$$A_1(j, T) = V(\phi_c) + j\phi_c - \frac{\log Z(\beta)}{\beta L^3} \tag{36}$$

in which we have also freely replaced $\bar{\phi}$ by ϕ_c in the logarithm of Z which itself is of order $\hbar$. Now by performing the Legendre transform (11), we find that the effective potential is

$$\begin{aligned}
V_1(\phi_c, T) &= V(\phi_c) - \frac{\log Z(\beta)}{\beta L^3} \\
&= V(\phi_c) + \int \frac{d^3k}{(2\pi)^3} \left[\frac{\omega_k}{2} + \frac{\log(1 - e^{-\beta\omega_k})}{\beta} \right]
\end{aligned} \tag{37}$$

with

$$\omega_k = \sqrt{k^2 + V''(\phi_c)}, \quad (38)$$

which is the usual result.

Classical potentials that induce spontaneous symmetry breaking have second derivatives that are negative between their inflection points. When the second derivative $V''(\phi_c)$ is negative, the frequency ω_k becomes complex for small enough k , and the loop expansion for the effective potential fails.

The preceding integral of ω_k over d^3k diverges. We may renormalize the effective potential by interpreting the classical potential $V(\phi)$ as containing counterterms $V_{ct}(\phi)$ of order $\hbar$ that are the same form as the terms of $V(\phi)$, apart from a constant term. By introducing a cut-off Λ and performing the integration, we find for the Helmholtz potential the expression

$$\begin{aligned} A_1(j, T) = & V(\phi_c) + j\phi_c + \frac{V''(\phi_c)^2}{64\pi^2} \left[\log \frac{V''(\phi_c)}{\mu^2} + \frac{1}{2} \right] \\ & + \frac{\Lambda^4}{16\pi^2} + \frac{\Lambda^2 V''(\phi_c)}{16\pi^2} + \frac{V''(\phi_c)^2}{64\pi^2} \log \frac{\mu^2}{4\Lambda^2} + V_{ct}(\phi_c) \\ & + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{-\rho(x)}) dx \end{aligned} \quad (39)$$

in which the renormalization point μ is arbitrary and $\rho(x) = \omega_k/T$ is the square root

$$\rho(x) = \sqrt{x^2 + V''(\phi_c)/T^2}. \quad (40)$$

Thus if we choose the counterterms minimally so that at ϕ_c they are

$$V_{ct}(\phi_c) = -\frac{\Lambda^4}{16\pi^2} - \frac{\Lambda^2 V''(\phi_c)}{16\pi^2} + \frac{V''(\phi_c)^2}{64\pi^2} \log \frac{4\Lambda^2}{\mu^2}, \quad (41)$$

then we obtain for the renormalized Helmholtz potential the formula

$$\begin{aligned} A_1(j, T) = & V(\phi_c) + j\phi_c + \frac{V''(\phi_c)^2}{64\pi^2} \left[\log \frac{V''(\phi_c)}{\mu^2} + \frac{1}{2} \right] \\ & + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{-\rho(x)}) dx, \end{aligned} \quad (42)$$

in which because of eq.(35) we may use either ϕ_c or $\bar{\phi}$ throughout. The effective potential is then given by the Legendre transform (11)

$$V_1(\phi_c, T) = A_1(j, T) - j\phi_c. \quad (43)$$

It will be useful in our discussion of the generalized effective potential to develop further the relation (34) between the mean value ϕ_c and the minimum $\bar{\phi}$ of the altered potential $V_j(\phi)$. By differentiating the extremal condition (26) with respect to j , we may find for the derivative of $\bar{\phi}$ the formula

$$\frac{\partial \bar{\phi}}{\partial j} = - \left[\frac{\partial^2 V(\bar{\phi})}{\partial \phi^2} \right]^{-1}. \quad (44)$$

It follows now from this formula and from eqs.(10), (34), and (42) that the mean value ϕ_c is to order $\hbar$

$$\phi_c = \bar{\phi} - \frac{V'''(\bar{\phi})}{32\pi^2} \left[\log \frac{V''(\bar{\phi})}{\mu^2} + 1 \right] - \frac{T^2 V'''(\bar{\phi})}{4\pi^2 V''(\bar{\phi})} \int_0^\infty \frac{x^2 dx}{\rho(x) (e^{\rho(x)} - 1)} \quad (45)$$

in which either ϕ_c or $\bar{\phi}$ may be used in the correction terms.

The archetypal example of a classical potential that exhibits spontaneous symmetry breaking is $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$. This potential has a positive second derivative $V''(\phi) = \lambda(3\phi^2 - \sigma^2)$ only for fields $|\phi|$ that are greater than $\sigma/\sqrt{3}$. For smaller $|\phi_c|$, the one-loop effective potential $V_1(\phi_c, T)$ is complex. According to eqs.(42) and (43), it is given by

$$\begin{aligned} V_1(\phi_c, T) = & \frac{\lambda}{4}(\phi_c^2 - \sigma^2)^2 + \frac{\lambda^2(3\phi_c^2 - \sigma^2)^2}{64\pi^2} \left[\log \left(\frac{\lambda(3\phi_c^2 - \sigma^2)}{\mu^2} \right) + \frac{1}{2} \right] \\ & + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{-\rho(x)}) dx \end{aligned} \quad (46)$$

where now $\rho(x)$ is

$$\rho(x) = \sqrt{x^2 + \lambda(3\phi_c^2 - \sigma^2)/T^2} \quad (47)$$

and μ is an arbitrary renormalization mass. To this expression one may add arbitrary, finite counterterms of the form

$$\lambda^2(A\phi_c^4 + B\phi_c^2 + C) \quad (48)$$

from the renormalization of the hamiltonian H . Due to the first logarithm $\log(\lambda(3\phi_c^2 - \sigma^2)/\mu^2)$, the effective potential $V_1(\phi_c, T)$ is complex for $|\phi_c| < \sigma/\sqrt{3}$, where it is not possible to quantize the approximate, altered theory.

The effective potential is the sum of a function of $\phi_c^2 - \sigma^2$ and a function of $3\phi_c^2 - \sigma^2$, and so cannot be convex. Its real part is concave, that is $\partial^2 \Re V_1(\phi_c, T)/\partial \phi_c^2 < 0$, for most of the interval $-\sigma/\sqrt{3} < \phi_c < \sigma/\sqrt{3}$ for small λ , low temperatures T , and reasonable renormalization.

For this example, the relation (45) between the mean value ϕ_c and the minimum $\bar{\phi}$ is

$$\phi_c = \bar{\phi} - \frac{3\lambda\bar{\phi}}{16\pi^2} \left[\log \frac{\lambda(3\bar{\phi}^2 - \sigma^2)}{\mu^2} + 1 \right] - \frac{3T^2\bar{\phi}}{2\pi^2(3\bar{\phi}^2 - \sigma^2)} \int_0^\infty \frac{x^2 dx}{\rho(x)(e^{\rho(x)} - 1)} \quad (49)$$

where $\rho(x)$ is given by eq.(47).

VI Where the Minima Lie

The multiple minima of symmetry-breaking classical potentials typically divide the space of fields into an inner region that includes the point $\phi = 0$ and an outer region that extends to infinite fields. The absolute minimum $\bar{\phi}$ of the altered potential $V_j(\phi) = V(\phi) + j\phi$ generally lies only in the outer region, which is not the region of interest in studies of the early universe.

Two examples may serve to illustrate this problem. For the classical potential $V(\phi) = \lambda(\phi^2 - \sigma^2)^2/4$, the current j and the global minimum $\bar{\phi}$ of the altered potential $V_j(\phi) = \lambda(\phi^2 - \sigma^2)^2/4 + j\phi$ have opposite signs. The absolute minimum $\bar{\phi}$ of $V_j(\phi)$ satisfies the extremal condition (26) which we may write in the form

$$\bar{\phi}^2 = \sigma^2 - j/(\lambda\bar{\phi}) \geq \sigma^2. \quad (50)$$

Thus in this example, the global minimum $\bar{\phi}$, which is ϕ_c to lowest order in $\hbar$,

always lies in the region $|\bar{\phi}| \geq \sigma$.

The second example is a theory of four scalar fields with a classical potential $V(|\phi|) = a + b|\phi|^2 + c|\phi|^4$ that has as its the absolute minimum the hypersphere $|\phi| = \sigma$. At the point $\phi_1 = |\phi|$ and $\phi_2 = \phi_3 = \phi_4 = 0$, the matrix $V_{ij}^{(2)}(\phi)$ of second derivatives of the potential is

$$V_{ij}^{(2)}(\phi) = \text{diag}(|\phi|V'', V', V', V') / |\phi|. \quad (51)$$

Only in the outer region, $|\phi| > \sigma$, are all the eigenvalues of this matrix positive. Thus the absolute minimum $\bar{\phi}$ must lie in this exterior region.

VII A More-Effective Potential

The reason for the complexity of the effective potential in models exhibiting spontaneous symmetry breaking is that the potential of the perturbed theory has the same second derivative $V_j''(\phi)$ as the original potential $V(\phi)$. Somewhat in the spirit of the composite-operator technique [25, 26], we may change the curvature of $V(\phi)$ by defining a more general Helmholtz potential $A(j, T; P)$ in which the linear probe $j\phi$ is replaced by a quadratic polynomial $jP(\phi)$

$$e^{-\beta L^3 A(j, T; P)} = \text{Tr} e^{-\beta(H + \int jP(\phi) d^3x)}. \quad (52)$$

On the one hand, it is clear that by this device we have not introduced any new divergences into the theory. On the other hand, it is also clear that the polynomial $P(\phi)$ is itself singular and requires regularization.

Now the derivative of the Helmholtz potential $A(j, T; P)$ with respect to the external current j is the mean value P_c

$$\frac{\partial A(j, T; P)}{\partial j} = P_c \quad (53)$$

defined as the mean value $\langle P(\phi) \rangle$ of the quadratic form $P(\phi)$

$$P_c = \frac{\text{Tr } P(\phi(x)) e^{-\beta(H + \int j P(\phi) d^3x)}}{\text{Tr } e^{-\beta(H + \int j P(\phi) d^3x)}}. \quad (54)$$

The classical field ϕ_c is still the mean value $\langle \phi \rangle$ of the quantum field ϕ

$$\phi_c = \frac{\text{Tr } \phi(x) e^{-\beta(H + \int j P(\phi) d^3x)}}{\text{Tr } e^{-\beta(H + \int j P(\phi) d^3x)}}. \quad (55)$$

We may now define a generalized effective potential $V(\phi_c, T; P)$ as the Legendre transform of the Helmholtz potential $A(j, T; P)$,

$$\begin{aligned} V(\phi_c, T; P) &= A(j, T; P) - j \frac{\partial A(j, T; P)}{\partial j} \\ &= A(j, T; P) - j P_c. \end{aligned} \quad (56)$$

This notation may be confusing. The variable that is conjugate to j is P_c ; so strictly speaking we probably should write $V(P_c, T; P)$ rather than $V(\phi_c, T; P)$. But all the potentials considered in this paper are actually and primarily functions of the external source j . And for a given perturbative ground state, the relationship between the source j and the mean value ϕ_c is one to one. Thus one may regard these potentials as functions of ϕ_c , which is the physically more-significant variable.

Like the conventional effective potential $V(\phi_c, T)$, the generalized effective potential at its minima describes the unperturbed system. For where the effective potential $V(\phi_c, T; P)$ is stationary, the external current j vanishes

$$\begin{aligned} 0 &= \frac{\partial V(\phi_c, T; P)}{\partial \phi_c} \\ &= \frac{\partial j}{\partial \phi_c} \left(\frac{\partial A(j, T; P)}{\partial j} - P_c \right) - j \frac{\partial P_c}{\partial \phi_c} \\ &= -j \frac{\partial P_c}{\partial \phi_c} \end{aligned} \quad (57)$$

unless P_c exceptionally should be independent of ϕ_c .

This more-effective potential $V(\phi_c, T; P)$ is real but not necessarily convex. Its second derivative contains two terms

$$\frac{\partial^2 V(\phi_c, T; P)}{\partial \phi_c^2} = -\frac{\partial j}{\partial \phi_c} \frac{\partial P_c}{\partial \phi_c} - j \frac{\partial^2 P_c}{\partial \phi_c^2} \quad (58)$$

and has no definite sign because the first term is typically positive while the second is typically negative.

We shall need the relation between the mean value ϕ_c and the external source j . To that end one may introduce the further-generalized Helmholtz potential $A(j_1, j_2, T; P)$ defined by the relation

$$e^{-\beta L^3 A(j_1, j_2, T; P)} = \text{Tr} e^{-\beta [H + \int (j_1 \phi + j_2 P) d^3 x]}. \quad (59)$$

Clearly when j_1 vanishes, this overly generalized Helmholtz potential reduces to the generalized Helmholtz potential:

$$A(0, j_2, T; P) = A(j_2, T; P). \quad (60)$$

The mean value ϕ_c for the more-effective potential is thus the partial derivative

$$\phi_c = \left. \frac{\partial A(j_1, j_2, T; P)}{\partial j_1} \right|_{j_1=0} \quad (61)$$

evaluated at $j_1 = 0$. On the other hand, the further-generalized Helmholtz potential $A(j_1, j_2, T; P)$ is the usual Helmholtz potential $A(j_1, T)$ for a theory in which the classical potential $V(\phi)$ has been shifted to $V(\phi) + j_2 P(\phi)$

$$A(j_1, j_2, T; P) = A(j_1, T)_{V+j_2 P}. \quad (62)$$

Thus by differentiating that potential $A(j_1, T)_{V+j_2 P}$ instead of $A(j_1, j_2, T; P)$ and by using eq.(10), we find that for the more-effective potential the mean value $\phi_c(j_2, T; P)$ is equal to the mean value $\phi_c(0, T)_{V+j_2 P}$ associated with the usual effective potential for a theory with a shifted potential $V + j_2 P$

$$\phi_c(j_2, T; P) = \phi_c(0, T)_{V+j_2 P} \quad (63)$$

evaluated at $j_1 = 0$.

By combining eqs.(60) and (62), we find that the generalized Helmholtz potential $A(j_2, T; P)$ is the usual Helmholtz potential $A(0, T)_{V+j_2P}$ for the shifted potential at vanishing j_1

$$A(j_2, T; P) = A(0, T)_{V+j_2P}. \quad (64)$$

In supersymmetric theories it may be appropriate to further generalize the perturbation P to a polynomial in both Fermi and Bose fields.

VIII The Meaning of Effective Potentials

The meaning of an effective potential is clearest at zero temperature. Since the perturbed hamiltonian $H + \int jP(\phi)d^3x$ is hermitian, it has eigenstates $|j\rangle$ with energy E_j

$$\left(H + \int jP(\phi)d^3x\right) |j\rangle = E_j |j\rangle. \quad (65)$$

Thus by its definition (52), the Helmholtz potential $A(j, T; P)$ in the limit $\beta \rightarrow \infty$ becomes the energy density $A(j, 0; P) = E_j/L^3$ of the eigenstate $|j\rangle$ of the altered hamiltonian $H + \int jP(\phi)d^3x$ with minimum energy E_j . And so by eq.(56), the effective potential

$$V(\phi_c, 0; P) = A(j, 0; P) - jP_c \quad (66)$$

is the mean value of the hamiltonian density in this state $|j\rangle$

$$V(\phi_c, 0; P) = \frac{\langle j|H|j\rangle}{L^3}. \quad (67)$$

And from eq.(57), it follows that the effective potential $V(\phi_c, 0; P)$ at its absolute minimum is in general the energy density of the ground state $|0\rangle$ of the unperturbed theory, *i.e.*, the energy density of the physical vacuum.

At finite temperatures, the potential $A(j, T; P)$ is the Helmholtz free-energy density of the mixture

$$\rho(j) = \frac{e^{-\beta[H + \int jP(\phi)d^3x]}}{\text{Tr } e^{-\beta[H + \int jP(\phi)d^3x]}} \quad (68)$$

associated with the altered hamiltonian density $[H + \int jP(\phi)d^3x]/L^3$. The finite-temperature effective potential $V(\phi_c, T; P)$ is the analog of the Gibbs free-energy density

$$V(\phi_c, T; P) = A(j, T; P) - j \frac{\partial A(j, T; P)}{\partial j} = A(j, T; P) - jP_c \quad (69)$$

of this mixture. By differentiating the definition (52) of the Helmholtz free-energy density $A(j, T; P)$ with respect to the temperature T , one may relate it to the perturbed energy density $u(j) = \text{Tr } \{\rho(j) [H + \int jP(\phi)d^3x]\} / L^3$ and the entropy density $s = -\text{Tr } \rho(j) \log \rho(j) / L^3$ of the mixture (68) by the equation

$$A(j, T; P) = u(j) - Ts. \quad (70)$$

It follows then from the relation (69) that the more-effective potential or Gibbs free-energy density $V(\phi_c, T; P)$ is related to the energy density $u = \text{Tr } \rho(j)H / L^3$ and entropy density s of the mixture $\rho(j)$ by the simpler equation

$$V(\phi_c, T; P) = u - Ts. \quad (71)$$

Unfortunately the mixture $\rho(j)$ given by eq.(68) coincides with the unperturbed physical mixture $\rho = e^{-\beta H} / \text{Tr } e^{-\beta H}$ only at the minima of $V(\phi_c, T; P)$ where the source j vanishes.

From this discussion it is clear that the choice of the polynomial $P(\phi)$ influences the effective potential $V(\phi_c, T; P)$ except at its various minima. In particular if one uses a quadratic polynomial P rather than a linear one, then one can avoid spurious complexities.

IX The One-Loop More-Effective Potential

Although the usual effective potential $V(\phi_c, T)$ is satisfactory for theories in which the classical potential $V(\phi)$ has positive curvature, it can become complex when $V''(\phi) < 0$, as in theories that exhibit spontaneous symmetry breaking. This section is concerned with the calculation of the one-loop, more-effective potential $V(\phi_c, T; P)$ for an arbitrary renormalizable classical potential $V(\phi)$. The computation will closely follow that of the usual effective potential.

If the curvature $V''(\phi)$ of the classical potential $V(\phi)$ is positive, then one may take $P(\phi) = \phi$ and the two effective potentials are identical. If the curvature $V''(\phi)$ of the classical potential $V(\phi)$ is negative for some range of ϕ , then the probe $P(\phi)$ should be quadratic. There are then two cases according to whether the classical potential has one or two minima.

If the potential $V(\phi)$ has a single minimum which we may call ϕ_1 , then we may take the probe $P(\phi)$ to be half the square of the distance from ϕ_1

$$P(\phi) \equiv P_1(\phi) = \frac{1}{2} (\phi - \phi_1)^2. \quad (72)$$

If the classical potential $V(\phi)$ has two minima, which we may call ϕ_1 and ϕ_2 , then it also has one local maximum ϕ_0 between them. In this case we may take the probe $P(\phi)$ to be half the square of the distance from the local maximum ϕ_0

$$P(\phi) \equiv P_2(\phi) = \frac{1}{2} (\phi - \phi_0)^2. \quad (73)$$

In both cases the first step is to find the minima $\bar{\phi}$ of the altered potential $V_j = V(\phi) + jP(\phi)$. These minima $\bar{\phi}$ are roots of the equation

$$0 = V'(\bar{\phi}) + jP'(\bar{\phi}). \quad (74)$$

Without any loss of generality, we may let the leading term of the potential $V(\phi)$ be $\lambda\phi^4/4$.

In the case of a unique minimum ϕ_1 , the derivative $V'(\phi)$ then will be the product of the three factors

$$V'(\phi) = \lambda (\phi - \phi_1) (\phi - z) (\phi - z^*). \quad (75)$$

So since the derivative of the probe $P_1(\phi)$ is

$$P'_1(\phi) = (\phi - \phi_1), \quad (76)$$

the minima $\bar{\phi}$ of the altered potential $V_j(\phi)$ are given by the quadratic equation

$$j = -\frac{V'(\bar{\phi})}{P'_1(\bar{\phi})} = -\lambda (\bar{\phi} - z) (\bar{\phi} - z^*) \leq 0. \quad (77)$$

We choose to compute the more-effective potential about the root $\bar{\phi}$ that is the absolute minimum of the altered potential $V_j(\phi)$.

When the potential $V(\phi)$ has two minima ϕ_1 and ϕ_2 , its derivative $V'(\phi)$ is the product of three factors

$$V'(\phi) = \lambda (\phi - \phi_0) (\phi - \phi_1) (\phi - \phi_2). \quad (78)$$

So since the derivative of the probe $P_2(\phi)$ is

$$P'_2(\phi) = (\phi - \phi_0), \quad (79)$$

the minima $\bar{\phi}$ of the altered potential $V_j(\phi) = V(\phi) + jP_2(\phi)$ are given by the quadratic equation

$$j = -\frac{V'(\bar{\phi})}{P'_2(\bar{\phi})} = -\lambda (\bar{\phi} - \phi_1) (\bar{\phi} - \phi_2). \quad (80)$$

The current j is positive for $\phi_1 < \bar{\phi} < \phi_2$. We choose to compute the more-effective potential about the root $\bar{\phi}$ that is the absolute minimum of the altered potential $V(\phi)$.

By its definition (52), the Helmolzt potential $A_0(j, T; P)$ in both cases and to lowest order in $\hbar$ is

$$A_0(j, T; P) = V(\bar{\phi}) + jP(\bar{\phi}). \quad (81)$$

To find the order- $\hbar$ correction to this result, we replace the altered potential $V + jP$ in the definition (52) of the generalized Helmholtz potential $A(j, T; P)$ by its truncated Taylor series

$$\begin{aligned} V(\phi) + jP(\phi) &\approx V(\bar{\phi}) + jP(\bar{\phi}) \\ &+ \frac{1}{2} \left(V''(\bar{\phi}) + jP''(\bar{\phi}) \right) (\phi - \bar{\phi})^2 \end{aligned} \quad (82)$$

and thereby reduce the problem to one that we have already solved (33). Thus including counterterms, we find

$$A_1(j, T; P) = V(\bar{\phi}) + V_{ct}(\bar{\phi}) + j \left(P(\bar{\phi}) + P_{ct}(\bar{\phi}) \right) - \frac{\log Z(\beta)}{\beta L^3}. \quad (83)$$

Here $V_{ct}(\bar{\phi})$ is the quartic polynomial (41) of counterterms that we used to renormalize the ordinary effective potential; $P_{ct}(\phi)$ is a quadratic polynomial in ϕ that we shall use to regularize the singular polynomial $P(\phi)$; and $P_{ct}(\bar{\phi})$ is that polynomial with the field ϕ replaced by its mean value $\bar{\phi}$. The squared mass in $Z(\beta)$ is positive and writable simply as

$$m^2 = V''(\bar{\phi}) + jP''(\bar{\phi}) = V''(\bar{\phi}) + j \geq 0 \quad (84)$$

because $\bar{\phi}$ is a minimum of $V_j(\phi)$ and because $P''(\bar{\phi}) = 1$ by construction (72–73). Thus the Helmholtz potential is

$$\begin{aligned} A_1(j, T; P) &= V(\bar{\phi}) + V_{ct}(\bar{\phi}) + j \left(P(\bar{\phi}) + P_{ct}(\bar{\phi}) \right) \\ &+ \int \frac{d^3 k}{(2\pi)^3} \left[\frac{\omega_k}{2} + \frac{\log(1 - e^{-\beta \omega_k})}{\beta} \right] \end{aligned} \quad (85)$$

in which now

$$\omega_k = \sqrt{k^2 + V''(\bar{\phi}) + j}. \quad (86)$$

If we again introduce a cut-off Λ and perform the integration, then we find

$$A_1(j, T; P) = V(\bar{\phi}) + jP(\bar{\phi}) + \frac{[V''(\bar{\phi}) + j]^2}{64\pi^2} \left[\log \frac{V''(\bar{\phi}) + j}{\mu^2} + \frac{1}{2} \right]$$

$$\begin{aligned}
& + \frac{\Lambda^4}{16\pi^2} + \frac{\Lambda^2[V''(\bar{\phi}) + j]}{16\pi^2} + \frac{[V''(\bar{\phi}) + j]^2}{64\pi^2} \log \frac{\mu^2}{4\Lambda^2} \\
& + V_{ct}(\bar{\phi}) + jP_{ct}(\bar{\phi}) + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{\rho(x)}) dx \quad (87)
\end{aligned}$$

in which μ is the renormalization point and $\rho(x)$ is the square root

$$\rho(x) = \sqrt{x^2 + [V''(\bar{\phi}) + j]}/T^2. \quad (88)$$

Using the same counterterms $V_{ct}(\bar{\phi})$ as the ones (41) we used for the usual effective potential, we obtain

$$\begin{aligned}
A_1(j, T; P) &= V(\bar{\phi}) + jP(\bar{\phi}) + \frac{[V''(\bar{\phi}) + j]^2}{64\pi^2} \left[\log \frac{V''(\bar{\phi}) + j}{\mu^2} + \frac{1}{2} \right] \\
&+ \frac{\Lambda^2 j}{16\pi^2} + \frac{2jV''(\bar{\phi}) + j^2}{64\pi^2} \log \frac{\mu^2}{4\Lambda^2} + jP_{ct}(\bar{\phi}) \\
&+ \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{\rho(x)}) dx. \quad (89)
\end{aligned}$$

All the terms on the right-hand side of this equation for $A_1(j, T; P)$ are functions of j , T , and the parameters of the theory. The same is true of $\bar{\phi}$. In particular the minimal choice of counterterms $P_{ct}(\phi)$ evaluated at $\bar{\phi}$ is

$$P_{ct}(\bar{\phi}) = -\frac{\Lambda^2}{16\pi^2} + \frac{2V''(\bar{\phi}) + j}{64\pi^2} \log \frac{4\Lambda^2}{\mu^2} \quad (90)$$

when expressed as a mixed function of j , $\bar{\phi}$, and the parameters of the theory. We shall now use the relations (77) and (80) that link j and $\bar{\phi}$ to write the counterterms $P_{ct}(\bar{\phi})$ as functions of $\bar{\phi}$ and the parameters of the theory without j . The counterterms $P_{ct}(\phi)$ will then be apparent.

In the case in which the classical potential $V(\phi)$ possesses a unique minimum ϕ_1 , we may identify the counterterms $P_{1,ct}(\phi)$ associated with the quadratic probe $P_1(\phi)$ by using the relation (77) between j and $\bar{\phi}$ to write the coefficient j of the logarithmically divergent term in (90) as $j = -\lambda |\bar{\phi} - z|^2$. The minimal

choice of counterterms $P_{1,ct}(\phi)$ then is

$$P_{1,ct}(\phi) = -\frac{\Lambda^2}{16\pi^2} + \frac{2V''(\phi) - \lambda|\phi - z|^2}{64\pi^2} \log \frac{4\Lambda^2}{\mu^2}. \quad (91)$$

These counterterms $P_{1,ct}(\phi)$ are of the same form as the probe $P_1(\phi)$, to wit a quadratic polynomial in the variable ϕ .

In the case in which the classical potential $V(\phi)$ possesses two minima ϕ_1 and ϕ_2 , we exploit the relationship (80) between j and $\bar{\phi}$ to write the same coefficient j of the logarithmically divergent term in (90) as $j = -\lambda(\bar{\phi} - \phi_1)(\bar{\phi} - \phi_2)$. The minimal choice of counterterms $P_{2,ct}(\phi)$ then is

$$P_{2,ct}(\phi) = -\frac{\Lambda^2}{16\pi^2} + \frac{2V''(\phi) - \lambda(\phi - \phi_1)(\phi - \phi_2)}{64\pi^2} \log \frac{4\Lambda^2}{\mu^2}. \quad (92)$$

Like the probe $P_2(\phi)$, these counterterms $P_{2,ct}(\phi)$ are a quadratic polynomial in the field ϕ .

The reason for the specific choices (72) and (73) of the probe polynomials $P_1(\phi)$ and $P_2(\phi)$ was so that their counterterms $P_{1,ct}(\phi)$ and $P_{2,ct}(\phi)$ would be of the same form as the probe polynomials themselves.

With the counterterms $P_{ct}(\phi)$, the Helmholtz potential is now

$$\begin{aligned} A_1(j, T; P) &= V(\bar{\phi}) + jP(\bar{\phi}) + \frac{[V''(\bar{\phi}) + j]^2}{64\pi^2} \left[\log \frac{V''(\bar{\phi}) + j}{\mu^2} + \frac{1}{2} \right] \\ &\quad + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{\rho(x)}) dx. \end{aligned} \quad (93)$$

This formula and its counterpart (42) with ϕ_c replaced by $\bar{\phi}$ are an example of the relation (64) between the generalized Helmholtz potential $A(j, T; P)$ and the usual Helmholtz potential $A(j, T)_{V+jP}$ for the theory with shifted potential $V + jP$.

To compute the more-effective potential $V_1(\bar{\phi}, T; P)$, one must exploit the relationship (77) or (80) between $\bar{\phi}$ and j and use eqs.(45) and (63) to relate the

mean value ϕ_c to the minimum $\bar{\phi}$:

$$\phi_c = \bar{\phi} - \frac{V'''(\bar{\phi})}{32\pi^2} \left[\log \frac{V''(\bar{\phi}) + j}{\mu^2} + 1 \right] - \frac{T^2 V'''(\bar{\phi})}{4\pi^2 (V''(\bar{\phi}) + j)} \int_0^\infty \frac{x^2 dx}{\rho(x) (e^{\rho(x)} - 1)} \quad (94)$$

in which either ϕ_c or $\bar{\phi}$ may be used in the correction terms and $\rho(x)$ is the square root (88). By again using the relationship (77) or (80) between $\bar{\phi}$ and j , one finally may perform the Legendre transform (56) and compute the more-effective potential $V_1(\bar{\phi}, T; P)$. We shall carry out this procedure explicitly for the classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$.

X An Example

This section contains a detailed computation of the generalized effective potential in the case of the classical potential

$$V(\phi) = \frac{\lambda}{4} (\phi^2 - \sigma^2)^2. \quad (95)$$

We have seen in Sec. V that if we use the linear polynomial $P(\phi) = \phi$, then the resulting effective potential is complex for $|\phi_c| < \sigma/3$. We shall find that by using the quadratic form $P(\phi) = \phi^2/2$, we may explore the whole region $|\phi_c| \leq \sigma$ with a more-effective potential $V(\phi_c, T; P)$ that remains real.

Since the potential (95) has indefinite curvature and two minima, the computation will follow the second of the two cases discussed in Sec. IX. The minima are

$$\phi_1 = -\sigma \quad \text{and} \quad \phi_2 = \sigma; \quad (96)$$

they are separated by a local maximum at $\phi = \phi_0 = 0$. So the probe is

$$P(\phi) = \frac{1}{2}\phi^2. \quad (97)$$

The minima $\bar{\phi}$ of the altered potential

$$V_j(\phi) = V(\phi) + \frac{j}{2}\phi^2 \quad (98)$$

are the roots of the quadratic equation

$$j = -\lambda(\bar{\phi}^2 - \sigma^2) \quad (99)$$

or

$$\bar{\phi}^2 = \sigma^2 - j/\lambda. \quad (100)$$

Let us choose to quantize about the positive root

$$\bar{\phi} = \bar{\phi}_+ = +\sqrt{\sigma^2 - (j/\lambda)}. \quad (101)$$

The mass associated with $\bar{\phi}$ is given by

$$m^2 = V_j''(\bar{\phi}) = 2\lambda\bar{\phi}^2 \geq 0. \quad (102)$$

Since the current j is related to the minimum $\bar{\phi}$ by (99), we may write this squared mass also as

$$m^2 = 2\lambda\sigma^2 - 2j. \quad (103)$$

To lowest order in $\hbar$, the Helmholtz potential $A_0(j, T; P)$ is

$$A_0(j, T; P) = V(\bar{\phi}) + j\frac{\bar{\phi}^2}{2} \quad (104)$$

in which $\bar{\phi}$ and j are related by $j = \lambda(\sigma^2 - \bar{\phi}^2)$. To this order the mean value ϕ_c and the minimum $\bar{\phi}$ are equal, and so the more-effective potential $V_0(\phi_c, T; P)$ is just the classical potential $V(\phi_c)$

$$V_0(\phi_c, T; P) = A_0(j, T; P) - j\frac{\partial A(j, T; P)}{\partial j} = \frac{\lambda}{4}(\phi_c^2 - \sigma^2)^2. \quad (105)$$

At the one-loop level, one finds by using eqs.(96) and (102) that the regularizing counterterms (92) evaluated at $\bar{\phi}$ are

$$P_{ct}(\bar{\phi}) = -\frac{\Lambda^2}{16\pi^2} + \frac{\lambda(5\bar{\phi}^2 - \sigma^2)}{64\pi^2} \log \frac{4\Lambda^2}{\mu^2}. \quad (106)$$

As a function of the external source j , the generalized Helmholtz potential $A_1(j, T; P)$ is then given by eq.(93) as

$$A_1(j, T; P) = \frac{j\sigma^2}{2} - \frac{j^2}{4\lambda} + \frac{(2\lambda\sigma^2 - 2j)^2}{64\pi^2} \left[\log \frac{2\lambda\sigma^2 - 2j}{\mu^2} + \frac{1}{2} \right] + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{\rho(x)}) dx \quad (107)$$

where now $\rho(x)$ is the square root

$$\rho(x) = \sqrt{x^2 + (2\lambda\sigma^2 - 2j)/T^2} = \sqrt{x^2 + 2\lambda\bar{\phi}^2/T^2}. \quad (108)$$

The generalized effective potential is defined (56) as the Legendre transform

$$V_1(\phi_c, T; P) = A(j, T; P) - j \frac{\partial A(j, T; P)}{\partial j}. \quad (109)$$

By performing the indicated differentiation with respect to j and by then expressing j as $\lambda(\sigma^2 - \bar{\phi}^2)$, we may write $V_1(\phi_c, T; P)$ as

$$V_1(\phi_c, T; P) = \frac{\lambda}{4}(\bar{\phi}^2 - \sigma^2)^2 + \frac{\lambda^2\bar{\phi}^2}{32\pi^2} \left[(4\sigma^2 - 2\bar{\phi}^2) \log \frac{2\lambda\bar{\phi}^2}{\mu^2} + 4\sigma^2 - 3\bar{\phi}^2 \right] + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \left[\log(1 - e^{-\rho(x)}) + \frac{\lambda(\sigma^2 - \bar{\phi}^2)}{T^2\rho(x)(e^{\rho(x)} - 1)} \right] dx \quad (110)$$

In the correction terms, which are of order $\hbar$, we may write indifferently $\bar{\phi}$ or ϕ_c . But in the first term $V(\bar{\phi})$ we must use eq.(94) to distinguish ϕ_c

$$\phi_c = \bar{\phi} - \frac{3\lambda\bar{\phi}}{16\pi^2} \left[\log \left(\frac{2\lambda\bar{\phi}^2}{\mu^2} \right) + 1 \right] - \frac{3T^2}{4\pi^2\bar{\phi}} \int_0^\infty \frac{x^2 dx}{\rho(x)(e^{\rho(x)} - 1)} \quad (111)$$

from $\bar{\phi}$. Since ϕ_c and $\bar{\phi}$ differ by terms of order $\hbar$, we need keep only the leading term

$$V(\bar{\phi}) \approx V(\phi_c) + \lambda\phi_c(\phi_c^2 - \sigma^2)(\bar{\phi} - \phi_c) \quad (112)$$

and may switch now to the variable ϕ_c throughout in the resulting formula for the one-loop, finite-temperature, generalized effective potential:

$$V_1(\phi_c, T; P) = \frac{\lambda}{4}(\phi_c^2 - \sigma^2)^2 + \frac{\lambda^2\phi_c^2}{32\pi^2} \left[(4\phi_c^2 - 2\sigma^2) \log \frac{2\lambda\phi_c^2}{\mu^2} + 3\phi_c^2 - 2\sigma^2 \right] + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \left[\log(1 - e^{-\rho(x)}) + \frac{\lambda(\phi_c^2 - \sigma^2)}{2T^2\rho(x)(e^{\rho(x)} - 1)} \right] dx \quad (113)$$

in which $\rho(x)$ is the square root

$$\rho(x) = \sqrt{x^2 + 2\lambda\phi_c^2/T^2}. \quad (114)$$

One may adopt a specific set of renormalization conditions by adding finite counterterms to the preceding formula. A sensible set of conditions is $V_1(\sigma, 0; P) = 0$, $V_1'(\sigma, 0; P) = 0$, and $V_1''(\sigma, 0; P) = m_H^2 = 2\lambda\sigma^2$. We may satisfy these conditions by adding the quartic polynomial

$$\frac{\lambda^2}{32\pi^2} (-8\phi^4 + 10\sigma^2\phi^2 - 3\sigma^4) \quad (115)$$

to the more-effective potential (113) and setting $\mu^2 = 2\lambda\sigma^2$. The resulting expression is

$$\begin{aligned} V_1(\phi_c, T; P) = & \frac{\lambda}{4}(\phi_c^2 - \sigma^2)^2 \\ & + \frac{\lambda^2}{32\pi^2} \left[(4\phi_c^4 - 2\sigma^2\phi_c^2) \log \frac{\phi_c^2}{\sigma^2} - 5\phi_c^4 + 8\sigma^2\phi_c^2 - 3\sigma^4 \right] \\ & + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \left[\log(1 - e^{-\rho(x)}) + \frac{\lambda(\phi_c^2 - \sigma^2)}{2T^2\rho(x)(e^{\rho(x)} - 1)} \right] dx \end{aligned} \quad (116)$$

with $\rho(x)$ given by eq.(114). If we set $\mu^2 = 2\lambda\sigma^2$ in the usual effective potential (46) and add to it these same counterterms, then it becomes

$$\begin{aligned} V_1(\phi_c, T) = & \frac{\lambda}{4}(\phi_c^2 - \sigma^2)^2 \\ & + \frac{\lambda^2}{64\pi^2} \left[(3\phi_c^2 - \sigma^2)^2 \log \left(\frac{3\phi_c^2 - \sigma^2}{2\sigma^2} \right) - \frac{23}{2}\phi_c^4 + 17\sigma^2\phi_c^2 - \frac{11}{2}\sigma^4 \right] \\ & + \frac{T^4}{2\pi^2} \int_0^\infty x^2 \log(1 - e^{-\rho(x)}) dx \end{aligned} \quad (117)$$

where now $\rho(x)$ is

$$\rho(x) = \sqrt{x^2 + \lambda(3\phi_c^2 - \sigma^2)/T^2}. \quad (118)$$

A numerical analysis of the formula (116) shows that the critical temperature runs from $T_C \approx 1.81\sigma$ for $\lambda = 0.1$ to $T_C \approx 1.7413\sigma$ for $\lambda = 1$. At higher

temperatures, the absolute minimum of $V_1(\phi_c, T; P)$ is at $\phi_c = 0$; at lower temperatures it is at $\phi_c > 0.62\sigma$ for $\lambda = 0.1$ and at $\phi_c > 0.69\sigma$ for $\lambda = 1$. The transition is weakly first order because at $T = T_C$ the barrier separating the two minima is slight. At the barrier temperature, which shifts from $T_B \approx 1.87\sigma$ for $\lambda = 0.1$ to $T_B \approx 1.865\sigma$ for $\lambda = 1$, this barrier disappears, and the field ϕ_c can roll classically from $\phi_c = \sigma$ to the absolute minimum at $\phi = 0$.

By differentiating eq.(116) with respect to ϕ_c at $\phi_c = 0$, one may show that at all positive temperatures, the derivative of the more-effective potential at $\phi_c = 0 + \epsilon$ is positive

$$\left. \frac{\partial V_1(\phi_c, T; P)}{\partial \phi_c} \right|_{\phi_c=0+\epsilon} > 0. \quad (119)$$

Thus the point $\phi_c = 0$ is a local minimum at all $T > 0$. The temperature T_2 at which this minimum disappears is therefore zero. In models of the early universe, inflation can occur if the field ϕ_c sticks at this local minimum.

The formula (117) for the usual effective potential $V_1(\phi_c, T)$ for $\lambda = 1$ is plotted (solid lines) in Fig. 1 at various temperatures. The region $0 \leq \phi_c < \sigma/\sqrt{3}$ is blank because the usual effective potential $V_1(\phi_c, T)$ is complex in this region. Clearly it is not possible to determine the critical temperature T_C or the barrier temperature T_B from the standard effective potential. The dotted curve represents the classical potential $V(\phi)$. The vertical axes are in units of $V(0) = \lambda\sigma^4/4$.

The formula (116) for the generalized effective potential $V_1(\phi_c, T; P)$ is plotted (solid lines) in Fig. 2 for $\lambda = 1$ at various temperatures including the critical temperature $T_C \approx 1.7413\sigma$, at which the minima at $\phi_c = 0$ and $\phi_c \approx 0.7\sigma$ are equally deep, and the barrier temperature $T_B \approx 1.865\sigma$, at which the barrier between these minima disappears. As in Fig. 1, the dotted curve represents the classical potential $V(\phi)$.

Figure 3 displays the generalized effective potential $V_1(\phi_c, T; P)$ as solid lines and the usual effective potential $V_1(\phi_c, T)$ as dashed lines for the same classical potential $V(\phi)$ (dots) with $\lambda = 1$. The more-effective potential tracks the standard effective potential so closely where the latter is real ($\phi_c \geq \sigma/\sqrt{3}$) that the dashes of $V_1(\phi_c, T)$ are visible only near $\phi_c = \sigma/\sqrt{3}$ and near $\phi_c = 1.4\sigma$. The generalized effective potential naturally extends the usual effective potential to $\phi_c = 0$.

By using the Haber–Weldon expansions of Bose–Einstein integrals [30], one may develop a high-temperature expansion [29] for the generalized effective potential (116)

$$\begin{aligned} V_1(\phi_c, T; P) = & -\frac{\pi^2}{90}T^4 + \frac{\lambda}{24}(3\phi_c^2 - \sigma^2)T^2 - \frac{\lambda^{3/2}}{12\sqrt{2}\pi}(7|\phi_c|^3 - 3\sigma^2|\phi_c|)T \\ & + \frac{\lambda^2}{16\pi^2}(2\phi_c^4 - \sigma^2\phi_c^2)\log\frac{T^2}{\sigma^2} + \frac{\lambda}{4}(\phi_c^2 - \sigma^2)^2 \\ & + \frac{\lambda^2}{32\pi^2}(f\phi_c^4 + s\phi_c^2 - 3\sigma^4) + \mathcal{O}\left(\frac{\sigma^2}{T^2}\right) \end{aligned} \quad (120)$$

in which $f = 4\log(8\pi^2/\lambda) - 8\gamma$, $s = 10 - 2\log(8\pi^2/\lambda) - 4\gamma$, and $\gamma \approx 0.57721566$. The high-temperature expansion [12] of the usual effective potential has the form

$$\begin{aligned} V_1(\phi_c, T) = & -\frac{\pi^2}{90}T^4 + \frac{\lambda}{24}(3\phi_c^2 - \sigma^2)T^2 - \frac{\lambda^{3/2}}{12\pi}(3\phi_c^2 - \sigma^2)^{3/2}T \\ & + \frac{\lambda^2}{64\pi^2}(3\phi_c^2 - \sigma^2)^2\log\frac{T^2}{\lambda^2(3\phi_c^2 - \sigma^2)^2} + \mathcal{O}(1). \end{aligned} \quad (121)$$

Although the third and fourth terms of this formula are imaginary for $|\phi_c| < \sigma/\sqrt{3}$, the first two terms, which are of order T^4 and T^2 , do agree with the first two terms of the high-temperature expansion (120) of the generalized effective potential (116). Thus the seemingly outrageous approximation, often made in work on the early universe, of keeping only these two, real terms of the high-temperature expansion (117) turns out to be a reasonable one after all.

Acknowledgements

I am grateful to D. Bailin, H. Barnum, S. Coleman, B. Grinstein, I. Hinchliffe, J. Jersák, S. Johnson, G. Keaton, E. Kolb, H. Leutwyler, S. Mandelstam, R. Matzner, J. March-Russell, F. Paige, R. Reeder, H. Stapp, G. Stephenson, P. Stevenson, T. Weiler, E. Weinberg, R. Xiu, and H. Zhang for helpful discussions. This research was supported by the U.S. Department of Energy under contracts DE-AC03-76SF00098 and DE-FG04-84ER40166. Some of this work was done at the Aspen Center for Physics.

References

- [1] W. Heisenberg and H. Euler, *Z. Phys.* 98 (1936) 714.
- [2] J. Schwinger, *Proc. Natl. Acad. Sci. USA* 37 (1951) 452 and *Phys. Rev.* 82 (1951) 664.
- [3] J. Goldstone, A. Salam, and S. Weinberg, *Phys. Rev.* 127 (1962) 965.
- [4] G. Jona-Lasinio, *Nuovo Cimento* 34 (1964) 1790.
- [5] J. Goldstone, *Nuovo Cimento* 19 (1961) 15; Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* 122 (1961) 345; 124 (1961) 246.
- [6] S. Coleman and E. Weinberg, *Phys. Rev. D* 7 (1973) 1888; S. Coleman, *Aspects of Symmetry* (Camb. Univ. Press, 1985), p. 132.
- [7] A. D. Linde, *JETP Letters* 23 (1976) 64.
- [8] S. Weinberg, *Phys. Rev. Lett.* 36 (1976) 294.
- [9] P. West, *Introduction to Supersymmetry and Supergravity* (World Scientific, 1990).

- [10] D. Kirzhnits and A. D. Linde, *Phys. Lett. B* 42 (1972) 471.
- [11] D. Kirzhnits and A. D. Linde, *JETP* 40 (1974) 628 and *Ann. Phys.* 101 (1976) 195.
- [12] L. Dolan and R. Jackiw, *Phys. Rev. D* 9 (1974) 3320.
- [13] S. Weinberg, *Phys. Rev. D* 9 (1974) 3357.
- [14] C. Bernard, *Phys. Rev. D* 9 (1974) 3312.
- [15] E. W. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley, 1990).
- [16] Y. Fujimoto, L. O’Raifeartaigh, and G. Parravicini, *Nucl. Phys. B* 212 (1983) 268; R. Haymaker and J. Perez-Mercader, *Phys. Rev. D* 15 (1983) 1948; R. J. Rivers, *Z. Phys. C* 22 (1984) 137; *Path-Integral Methods in Quantum Field Theory* (Cambridge, 1987), pp. 235–272; C. M. Bender and F. Cooper, *Nucl. Phys. B* 224 (1983) 403; F. Cooper and B. Freedman, *Nucl. Phys. B* 239 (1984) 459.
- [17] K. Symanzik, *Commun. Math. Phys.* 16 (1970) 48; J. Iliopoulos, C. Itzykson, and A. Martin, *Rev. Mod. Phys.* 47 (1975) 165.
- [18] E. J. Weinberg and A. Wu, *Phys. Rev. D* 36 (1987) 2474.
- [19] S. J. Chang, *Phys. Rev. D* 12 (1975) 1071.
- [20] T. Barnes and G. I. Ghandour, *Phys. Rev. D* 22 (1980) 924.
- [21] P. M. Stevenson, *Phys. Rev. D* 30 (1984) 1712; 32 (1985) 1389; *Z. Phys.* C35 (1987) 467.
- [22] R. Fukuda and E. Kyriakopoulos, *Nucl. Phys. B* 85 (1975) 354; R. Fukuda, *Prog. Theor. Phys.* 56 (1976) 258.

- [23] L. O’Raifeartaigh, A. Wipf, and H. Yoneyama, *Nucl. Phys. B* 271 (1986) 653.
- [24] A. Ringwald and C. Wetterich, *Nucl. Phys.* B334 (1990) 506.
- [25] J. M. Cornwall, R. Jackiw, and E. Tomboulis, *Phys. Rev. D* 10 (1974) 2428.
- [26] S. W. Hawking and I. G. Moss, *Nucl. Phys. B* 224 (1983) 180.
- [27] I. D. Lawrie, *Nucl. Phys.* B301 (1988) 685.
- [28] G. W. Anderson and L. J. Hall, *Phys. Rev.* D45 (1992) 2685.
- [29] K. Cahill and T. Weiler, work in progress.
- [30] H. E. Haber and H. A. Weldon, *J. Math. Phys.* 23 (1982) 1852.




Figure 1: For the classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$ with $\lambda = 1$ (dots), the usual effective potential $V_1(\phi_c, T)$ (solid lines) is plotted where it is real $\phi_c > \sigma/\sqrt{3}$ in units of $V(0) = \lambda\sigma^4/4$ at various temperatures including the critical temperature $T_C \approx 1.741\sigma$ and the barrier temperature $T_B \approx 1.865\sigma$.




Figure 2: For the classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$ with $\lambda = 1$ (dots), the generalized effective potential $V_1(\phi_c, T; P)$ is plotted (solid lines) in units of $V(0) = \lambda\sigma^4/4$ at various temperatures including the critical temperature $T_C \approx 1.741\sigma$ at which the minima at $\phi_c = 0$ and $\phi_c \approx 0.7\sigma$ are equally deep and the barrier temperature $T_B \approx 1.865\sigma$ at which the barrier between these minima disappears.




Figure 3: For the classical potential $V(\phi) = (\lambda/4)(\phi^2 - \sigma^2)^2$ with $\lambda = 1$ (dots), the usual effective potential $V_1(\phi_c, T)$ (dashes) and the more-effective potential $V_1(\phi_c, T : P)$ (solid lines) are plotted in units of $V(0) = \lambda\sigma^4/4$ at various temperatures. The more-effective potential tracks the standard effective potential very closely where the latter is real ($\phi_c > \sigma/\sqrt{3}$) and naturally extends it to $\phi_c = 0$.